

# First restriction and genetic mapping of the genomic DNA of urease-positive thermophilic campylobacters (UPTC), and small restriction fragment sequencing

T. ARITOMI<sup>1</sup>, T. SEKIZUKA<sup>2</sup>, R. IMAMAKI<sup>1</sup>, O. MURAYAMA<sup>1</sup>, B. C. MILLAR<sup>3</sup>, J. E. MOORE<sup>3</sup> and M. MATSUDA<sup>4</sup>

<sup>1</sup>Laboratory of Molecular Biology, School of Environmental Health Sciences, Azabu University, Sagamihara, Japan; and <sup>2</sup>Department of Bacteriology, Northern Ireland Public Health Laboratory, Belfast City Hospital, Belfast BT9 7AD, Northern Ireland, UK

An incorrect reference (ref. 19) was cited when this paper was published originally (*Br J Biomed Sci* 2006; **63**: 63–7). The reference section from this paper is reproduced here and includes the correct citation for reference 19.

## References

- Skirrow MB, Benjamin J. Differentiation of enteropathogenic *Campylobacter*. *J Clin Pathol* 1980; **33**: 1122.
- Benjamin J, Leaper S, Owen RJ, Skirrow MB. Description of *Campylobacter laridis*, a new species comprising the nalidixic acid-resistant thermophilic *Campylobacter* (NARTC) group. *Curr Microbiol* 1983; **8**: 231–8.
- Bolton FJ, Holt AV, Hutchinson DN. Urease-positive thermophilic campylobacters. *Lancet* 1985; **i**: 1217–8.
- Owen RJ, Costas M, Sloss L, Bolton J. Numerical analysis of electrophoretic protein patterns of *Campylobacter laridis* and allied thermophilic campylobacters from the natural environment. *J Appl Bacteriol* 1988; **65**: 69–78.
- Megraud F, Chevrier D, Desplaces N, Sedallian A, Guesdon J L. Urease-positive thermophilic *Campylobacter* (*Campylobacter laridis* variant) isolated from an appendix and from human faeces. *J Clin Microbiol* 1988; **26**: 1050–1.
- Bezian MC, Ribou G, Baberis-Giletti C, Megraud F. Isolation of urease positive thermophilic variant of *Campylobacter lari* from a patient with urinary tract infection. *Eur J Clin Microbiol Infect Dis* 1990; **9**: 895–7.
- Wilson IG, Moore JE. Presence of *Salmonella* spp. and *Campylobacter* spp. in shellfish. *Epidemiol Infect* 1996; **116**: 147–53.
- Endtz HP, Vliegthart JS, Vandamme P *et al.* Genotypic diversity of *Campylobacter lari* isolated from mussels and oysters in The Netherlands. *Int J Food Microbiol* 1997; **34**: 79–88.
- Kaneko A, Matsuda M, Miyajima M, Moore JE, Murphy PG. Urease-positive thermophilic strains of *Campylobacter* isolated from seagulls (*Larus* spp.). *Lett Appl Microbiol* 1999; **29**: 7–9.
- Matsuda M, Kaneko A, Stanley T *et al.* Characterisation of urease-positive thermophilic *Campylobacter* subspecies by multilocus enzyme electrophoresis typing. *Appl Environ Microbiol* 2003; **69**: 3308–10.
- Matsuda M, Kaneko A, Fukuyama M *et al.* First finding of urease-positive thermophilic strains of *Campylobacter* in river water in the Far East, namely, in Japan, and their phenotypic and genotypic characterization. *J Appl Bacteriol* 1996; **81**: 608–12.
- Matsuda M, Shibuya T, Itoh Y *et al.* First isolation of urease-positive thermophilic *Campylobacter* (UPTC) from crows (*Corvus leuclantii*) in Japan. *Int J Hyg Environ Health* 2002; **250**: 321–4.
- Sambrook J, Fritsch E F, Maniatis T. *Molecular cloning, a laboratory manual* (2nd edn). New York: Cold Spring Harbor Laboratory Press, 1989.
- Newnham E, Chang N, Taylor DE. Expanded genomic map of *Campylobacter jejuni* UA580 and localization of 23S ribosomal rRNA genes by I-CeuI restriction endonuclease digestion. *FEMS Microbiol Lett* 1996; **142**: 223–9.
- Al Rashid ST, Dakuna I, Louie H *et al.* Identification of *Campylobacter jejuni*, *C. coli*, *C. lari*, *C. upsaliensis*, *Arcobacter butzleri*, and *A. butzleri*-like species based on the *glyA* gene. *J Clin Microbiol* 2000; **38**: 1488–94.
- Miyajima M, Matsuda M, Haga S, Kagawa S, Millar BC, Moore JE. Cloning and sequencing of 16S rDNA and 16S-23S rDNA internal spacer region (ISR) from urease-positive thermophilic *Campylobacter* (UPTC). *Lett Appl Microbiol* 2002; **34**: 287–9.
- Sekizuka T, Gondo T, Murayama O *et al.* Molecular cloning, nucleotide sequencing and characterization of the flagellin gene from isolates of urease-positive thermophilic *Campylobacter*. *Res Microbiol* 2004; **155**: 185–91.
- Honda M, Matsushita S, Murayama O, Millar BC, Moore JE, Matsuda M. Cloning and sequence analysis of the *recA* gene in urease-positive thermophilic campylobacters (UPTC). *Br J Biomed Sci* 2002; **59**: 166–9.
- Usui K, Iida H, Ueno H *et al.* Genetic heterogeneity of urease gene loci in urease-positive thermophilic *Campylobacter* (UPTC). *Int J Hyg Environ Health* 2006. In press (Epub ahead of print).
- Parkhill J, Wren BW, Mungall K *et al.* The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature* 2000; **403**: 665–8.
- Regonesi ME, Briani F, Ghetta A *et al.* A mutation in polynucleotide phosphorylase from *Escherichia coli* impairing RNA binding and degradosome stability. *Nucleic Acids Res* 2004; **32**: 1006–17.
- Fouts DE, Mongodin EF, Mandrell RE *et al.* Major structural differences and novel potential virulence mechanisms from the genomes of multiple *Campylobacter* species. *PLoS Biol* 2005; **3**: e15.
- Sekizuka T, Gondo T, Murayama O, Moore JE, Millar BC, Matsuda M. *flaA*-like sequences containing internal termination codons (TAG) in urease-positive thermophilic *Campylobacter* isolated in Japan. *Lett Appl Microbiol* 2002; **35**: 185–9.

Correspondence to: Dr. Motoo Matsuda  
Email, matsuda@azabu-u.ac.jp